Feeding the Follicle: A Functional Nutrition Framework for Hair Loss in Women

Julie Olson, BCHN®, CN, CFMP, CGP, BSc

Abstract

Hair loss in women is routinely attributed to hormones, genetics, or stress, yet an underrecognized and highly correctable driver often underlies all three: nutritional insufficiency. This article presents a functional-nutrition framework for evaluating and addressing nutrient status in women presenting with diffuse or patterned hair loss, synthesizing peer-reviewed literature on protein, iron, the B-vitamin complex, zinc, selenium, the fat-soluble vitamins, and omega-3 fatty acids (Almohanna et al., 2019; Gomes et al., 2025). Particular attention is given to the gap between conventional laboratory reference ranges and the functional thresholds required for active follicle growth, a gap that leaves many women told their labs are "normal" even as their hair continues to thin. The literature reviewed spans systematic reviews, meta-analyses, and clinical trials published predominantly within the past five to eight years, supplemented by a small number of foundational studies that first established key mechanisms and have not been superseded. The article closes with a clinical decision framework, test, don’t guess, for nutrition professionals evaluating hair loss in practice, along with guidance on where food-based intervention, targeted supplementation, and gut-absorption support each belong in a comprehensive plan.

Introduction

Lisa had done everything right. A whole-foods diet, mostly organic. Lean protein at every meal. A supplement stack her naturopath had helped her build. Six months in, she was still watching her hair thin. "I don’t understand," she said at our first session. "I’m eating better than I ever have in my life."

She was right. She was eating better. But her ferritin was 18 ng/mL, technically within the conventional reference range yet functionally insufficient for the follicles that needed it. Her vitamin D was 24 ng/mL. Her zinc was at the low end of "normal." And her gut was not absorbing what she was eating at the cellular level where it counted.

Lisa’s case is not unusual. It is, in fact, the pattern most consistently observed in clinical practice among women presenting for hair loss: an intact diet, a reasonable supplement routine, and labs that appear unremarkable, alongside progressive shedding that conventional workups cannot explain. A 2025 systematic review of dietary factors and

hair health, drawing on seventeen studies and more than sixty-one thousand participants, found that higher serum vitamin D and adequate iron status were consistently associated with better hair outcomes, while higher intake of alcohol and sugar-sweetened beverages was linked to increased hair loss risk (Gomes et al., 2025). This finding aligns with a broader body of review-level evidence establishing vitamins and minerals as legitimate, modifiable contributors to hair loss rather than a fringe concern (Almohanna et al., 2019). The evidence base linking nutrition to hair health has matured considerably. What has not kept pace, in most clinical settings, is the translation of that evidence into how hair loss is screened and treated.

Nutrition is not a substitute for the broader root-cause work that hair loss usually requires, including hormonal, gut, and stress-related evaluation. But deficiency is a direct upstream cause of hair loss, and until it is identified and corrected, no amount of work in those other domains will produce the results that become possible once nutritional status is properly aligned. This article outlines which nutrients the hair follicle requires, explains why deficiency is more common than standard testing reveals, and details what a nutrition professional needs to know to build an evidence-informed plan for a client whose hair loss has a nutritional component.

Methodology

This article synthesizes peer-reviewed literature on nutrition and hair loss identified through PubMed/MEDLINE and Google Scholar searches combining hair loss, alopecia, telogen effluvium, and androgenetic alopecia with deficiency, micronutrient, and supplementation terms. Priority was given to systematic reviews, meta-analyses, and controlled clinical trials published within the past five to eight years. A small number of older studies were retained when they established a mechanism or clinical threshold that has not since been superseded or replicated at the same scale; these are identified as such in the reference list. Findings were synthesized through a functional-nutrition clinical lens developed in practice with women experiencing hair loss, organized around a single governing principle: confirm deficiency through functional laboratory testing before treating for it, and address the gut's capacity to absorb nutrients before assuming that a supplement protocol, however well-designed, will resolve what the labs show.

The Absorption Gap

Before turning to individual nutrients, one foundational reality deserves to be stated directly. The most common nutritional pattern in women with hair loss is not simply "not eating enough of the right things." It is an absorption gap: the nutrients are present in the diet and often in the supplement routine, but the gut is not delivering them to the

follicle at the concentration required for healthy growth. Gut dysfunction, including reduced stomach acid, intestinal inflammation, and dysbiosis, causes malabsorption even in diets that are nutritionally excellent on paper. Emerging research on the gut-follicle axis is beginning to map the biological pathways behind this pattern (Liu & Liu, 2023).

This does not make dietary choices irrelevant. Food quality, nutrient density, and eating patterns matter enormously (Almohanna et al., 2019). But it means that functional testing, gut restoration, and targeted supplementation must accompany dietary improvement for the full clinical picture to resolve. The guiding principle throughout this article: test, don't guess. Supplement to correct a confirmed deficiency, not as a precaution against a hypothetical one. And address the gut's absorptive capacity before expecting supplementation to work, because without a functioning gut lining, a client may be investing in nutrients her body cannot use.

Protein: The Keratin Foundation

Hair is primarily composed of keratin, a structural protein. Without sufficient dietary protein, the body enters a biological triage state in which protein delivery is prioritized for organs and systems essential for survival. Because hair is non-critical in the body's hierarchy, it is cut off from the supply chain. This mechanism was first documented nearly five decades ago in women on severely calorie- and protein-restricted diets, who developed diffuse shedding that resolved once adequate protein intake resumed. It remains one of the more consistently reproduced findings in the nutrition-hair literature (Goette & Odom, 1976). Subsequent review-level evidence has reinforced protein adequacy as a foundational, if often overlooked, variable in hair loss evaluation (Almohanna et al., 2019). Protein deficiency-induced hair loss typically presents as diffuse shedding rather than patterned loss, a generalized thinning that can develop within two to three months of inadequate intake and accelerate rapidly once it begins.

The relevant clinical question is not whether a client eats protein, but whether she consumes adequate complete protein: sources that provide all nine essential amino acids the body cannot synthesize on its own. Animal proteins (eggs, fatty fish, lean meat, dairy) are complete by definition; most plant proteins are not, with soy, quinoa, and hemp seed as notable exceptions. This is not an argument against plant-based eating, but an argument for recognizing that plant-protein combinations require intentional planning when they serve as a primary protein source.

A practical target for women supporting hair health is 1.0 to 1.2 grams of protein per kilogram of body weight daily, at the higher end of general population

recommendations, given the follicle's metabolic demands and the rapid cell turnover required to sustain active growth phases. In practical terms, this typically translates to roughly 60 to 80 grams of high-quality protein daily, distributed across meals in portions of about 20 grams each, such as a piece of chicken, fish, or an equivalent protein source about the size of a palm.

Iron: The Most Common and Most Overlooked Deficiency

Iron deficiency is widely regarded as the most common nutritional cause of hair loss in women and remains systematically undertested in standard clinical workups (Almohanna et al., 2019). A 2025 case-control study comparing one hundred women with hair loss to age-matched controls found significantly lower hemoglobin, serum iron, and ferritin in the hair loss group, along with lower selenium, zinc, copper, calcium, vitamin B12, and vitamin D. Ferritin was identified as one of several independent predictors of hair loss on multivariable analysis (Al-Fawaeir & Al-Odat, 2025). A separate 2025 study of women with telogen effluvium found serum ferritin nearly half that of healthy controls (24.3 ng/mL versus 44.8 ng/mL). Clinical iron deficiency was present in over a quarter of affected women and absent in the control group entirely (Rangarajan, 2025). A 2024 biochemical survey of chronic telogen effluvium patients likewise included ferritin among the panel of markers distinguishing affected women from controls, reinforcing iron status as a recurring finding across independent research groups rather than an isolated result (Durusu Turkoglu et al., 2024).

The core clinical problem is rarely serum iron; it is ferritin, the protein that stores iron intracellularly. Conventional labs often flag ferritin as "normal" at levels above 10 to 12 ng/mL. For hair follicle function, the functional threshold is considerably higher, and this is precisely the gap that leaves women told their iron is "fine" even as their hair continues to thin. It is also exactly why functional laboratory testing is not optional in a hair loss workup: it is the only way to see where ferritin actually sits relative to what the follicle needs, rather than relative to a reference range built around anemia prevention.

Iron from animal sources, heme iron, found in red meat, poultry, and shellfish, is absorbed at roughly two to three times the rate of plant-source (non-heme) iron (Almohanna et al., 2019). For clients who are primarily plant-based, pairing iron-rich plant foods with a source of vitamin C at the same meal substantially improves non-heme iron absorption. Healing the gut lining removes a significant additional barrier to absorption, regardless of dietary source.

The B-Vitamin Complex

Three B vitamins are particularly relevant to hair health, and biotin has accumulated a mythology worth examining from both directions. Conventional medicine has often dismissed biotin as clinically irrelevant to hair loss in the absence of an obvious cause, such as raw egg white consumption, which blocks biotin absorption, or specific medications known to deplete it. A 2026 systematic review of biotin supplementation trials for alopecia found that biotin monotherapy did not produce consistent, objectively measured hair growth benefits, and that improvements reported in combination-ingredient products could not be attributed specifically to biotin (Moltó-Balado et al., 2026). At the same time, biotin status is measured inconsistently across the hair loss population, and a 2024 biochemical analysis of women with chronic telogen effluvium assessed biotin alongside ferritin, vitamin B12, vitamin D, zinc, copper, and selenium, underscoring that biotin deficiency, where it exists, rarely occurs in isolation (Durusu Turkoglu et al., 2024). The correct clinical position is not to dismiss biotin or megadose it reflexively, but to test, confirm a true deficiency, and correct it. Gut dysbiosis can also produce biotin insufficiency from the inside out, since gut bacteria synthesize biotin, and a disrupted microbial ecosystem can create deficiency even without poor dietary intake.

Folate (B9) directly supports the rapid cell division that hair follicle cells, among the fastest-dividing cells in the human body, depend on (Almohanna et al., 2019). Deficiency impairs this proliferation, resulting in slower, weaker growth. Clients on hormonal contraceptives or methotrexate may experience folate depletion, a known pharmacological side effect. This connection rarely surfaces in a standard hair-loss workup but carries real clinical significance.

Vitamin B12 drives red blood cell production and oxygen delivery, and hair follicles are among the more oxygen-demanding tissues in the body; insufficient B12 limits the oxygen supply required to sustain active growth phases (Almohanna et al., 2019). A 2025 case-control study similarly found significantly lower vitamin B12 levels among women with hair loss compared to healthy controls, identifying it as an independent predictor alongside ferritin, folate, and vitamin D (Al-Fawaeir & Al-Odat, 2025). Clients following plant-based diets and women over fifty, whose intrinsic factor production naturally declines and reduces B12 absorption, carry elevated risk. Standard serum B12 testing has a known false-negative problem, since levels at the low end of "normal" can reflect functional deficiency at the cellular level. Therefore, methylmalonic acid and homocysteine are useful adjunct markers of B12 sufficiency.

Zinc and Selenium: The Minerals of Precision

Zinc supports protein synthesis pathways that build keratin, anti-inflammatory regulation that protects follicles from chronic immune activation, and enzymatic processes involved in thyroid hormone conversion (Almohanna et al., 2019). A 2022 systematic review and meta-analysis found significantly lower serum zinc levels in patients with androgenetic alopecia compared with controls, supporting zinc's relevance to this pattern of hair loss (Chang & Chang, 2022). A 2024 biochemical study of chronic telogen effluvium found zinc to be the single nutrient most significantly associated with hair shedding among a full panel of nutrients tested (Durusu Turkoglu et al., 2024). That said, the evidence is not uniform: a 2025 cross-sectional study of nearly twenty-four thousand patients found only a small difference in median zinc levels between those with and without hair loss diagnoses, with both groups falling within the normal physiological range, and concluded that routine zinc testing is not, by itself, a useful screen for hair loss (Liran et al., 2025). Taken together, these findings support zinc as one factor worth evaluating within a broader nutrient panel rather than a stand-alone diagnostic marker, and they reinforce why self-prescribing zinc without confirmed testing is inadvisable: excess zinc competes with copper absorption and can disrupt the mineral balance that supports healthy follicle function.

Selenium supports the conversion of thyroid hormone from its inactive form (T4) to its active form (T3), directly linking it to the hormonal dimension of hair loss (Almohanna et al., 2019). Brazil nuts are an exceptionally concentrated dietary source; two to three per day meet the daily selenium requirement for most women. As with zinc, selenium is a nutrient for which both deficiency and excess create problems, and supplementing without confirmed deficiency is unnecessary and, at higher doses, potentially harmful.

The Fat-Soluble Vitamins

Vitamin D may be among the most significant and most commonly deficient nutrients in women with hair loss. Hair follicle cells carry vitamin D receptors, and a 2024 systematic review and meta-analysis of eighty-one studies involving more than fifteen thousand patients with alopecia found that vitamin D deficiency was substantially more prevalent in affected individuals than in controls, most notably in alopecia areata (odds ratio 2.84) and female pattern hair loss (odds ratio 5.24) (Yongpisarn et al., 2024). Two independent 2024 and 2025 case-control studies found the same pattern in women with hair loss and telogen effluvium, with vitamin D emerging as one of several independent predictors of shedding (Al-Fawaeir & Al-Odat, 2025; Durusu Turkoglu et al., 2024). Functional practitioners typically target serum 25-hydroxyvitamin D above 50 ng/mL

for optimal hair support, a level most American women do not reach without supplementation, given limited sun exposure at northern latitudes and the scarcity of high-concentration dietary sources. Testing before supplementing matters here too: vitamin D toxicity from oversupplementation, while less common than deficiency, is a real clinical concern.

Vitamin C serves two critical functions in hair nutrition. It is an essential cofactor for collagen synthesis, and collagen forms part of the dermal papilla structure that anchors the follicle in the scalp, and it substantially enhances iron absorption from plant sources when consumed at the same meal (Almohanna et al., 2019). As a potent antioxidant, vitamin C also neutralizes oxidative stress generated by gut dysfunction and chronic inflammation, protecting follicles from cellular damage that prematurely shortens the growth cycle.

Vitamin A is essential for sebum production, the natural oil that lubricates the scalp and maintains the follicular environment, yet it is one of the clearest examples in nutritional medicine of a nutrient for which more is not better (Almohanna et al., 2019). Both deficiency and excess cause hair loss. Preclinical research on dietary vitamin A's effects on the hair cycle found that excessive intake arrested follicles in a prolonged resting phase, while inadequate intake produced the same result through a different mechanism, pointing to a narrow physiological window in either direction (Suo et al., 2021). Self-prescribing high-dose vitamin A or retinol without confirmed blood levels is a common and often overlooked nutritional error, frequently made by clients who have read that "vitamin A is good for skin" without understanding how narrow its therapeutic window actually is.

Vitamin E provides antioxidant protection at the scalp level and supports microvascular function that delivers nutrients to the follicle. Tocotrienol forms of vitamin E, less common in commercial supplements than tocopherol forms, have shown specific benefit for hair density in clinical research: a randomized, placebo-controlled trial found that oral tocotrienol supplementation produced a significant increase in hair count over eight months in volunteers with hair loss. This finding remains, more than a decade later, one of the few controlled trials isolating this specific form of vitamin E (Beoy et al., 2010).

Omega-3 Fatty Acids and the Gut-Inflammation Connection

Omega-3 fatty acids, found in fatty fish, flaxseed, chia seeds, and walnuts, act as anti-inflammatory agents at the cellular level. For hair, their primary value lies in reducing scalp inflammation that shortens growth phases and impairs follicle function, a pathway increasingly implicated in the gut-hair connection. A growing body of research on the

gut microbiome's role in alopecia areata has identified specific bacterial signatures associated with disease status and documented gut-derived short-chain fatty acids as modulators of the inflammatory and immune signaling pathways that govern the hair growth cycle, reinforcing why an anti-inflammatory nutrition strategy and gut restoration work belong together rather than as separate interventions (Liu & Liu, 2023). The standard Western diet is dramatically skewed toward the pro-inflammatory omega-6 end of the fat spectrum relative to omega-3, a pattern consistent with the broader dietary-risk findings described earlier (Gomes et al., 2025). Correcting this ratio through both dietary shifts and targeted supplementation is one of the most impactful, lowest-risk nutritional adjustments most women can make, regardless of where the rest of their nutritional picture stands. In practice, introducing a better-balanced intake of these fats has often produced a visible improvement in clients with dry, lifeless hair, even before other interventions take full effect.

Supplements Versus Food: A Clinical Framework

Both the supplement industry's messaging and conventional medicine's reflexive dismissal of nutritional interventions miss the actual clinical picture. A 2023 systematic review in JAMA Dermatology examined thirty studies of nutritional supplements for hair loss and found that interventions with the highest-quality supporting evidence showed potential benefit, that no diet-based interventional studies met inclusion criteria (underscoring how little controlled research exists on whole-food dietary patterns specifically), and that adverse effects across the reviewed therapies were rare and mild (Drake et al., 2023). A more recent meta-analysis reached a similarly measured conclusion, finding a positive signal for nutraceutical supplementation in non-scarring alopecia while cautioning that study heterogeneity limits how confidently those findings generalize to individual clients (Ahmed et al., 2025). A controlled trial of a combination oral supplement containing amino acids, iron, selenium, and marine hydrolyzed collagen found measurable improvement in hair loss outcomes over three months, illustrating the kind of multi-nutrient formulation that performs better in trials than single isolated nutrients typically do (Milani & Colombo, 2023).

Whole foods remain the foundation. The nutrient cofactors, fiber, and phytonutrients in food work synergistically in ways that isolated supplement compounds cannot replicate (Gomes et al., 2025). A diet that includes fatty fish, leafy greens, eggs, seeds, and colorful vegetables provides not only nutrients but also the cofactors that help those nutrients function, and it feeds the gut bacteria that produce additional nutrients internally.

But supplementation is essential when a confirmed deficiency exists. Food alone cannot reliably correct severe iron deficiency, vitamin D insufficiency in clients at northern latitudes with minimal sun exposure, or B12 deficiency in those following plant-based diets. In these cases, targeted supplementation under clinical guidance is not optional; it is required to restore normal function. The governing principle for practitioners: test first, then supplement to correct what testing reveals, not as insurance against a hypothetical deficiency, and not before the gut's absorptive capacity has been addressed, because supplementing into a system that cannot utilize what is provided is an expensive exercise in futility.

Emerging and Adjunctive Nutrients

Clients frequently ask about a long list of additional ingredients marketed for hair health: curcumin, ginkgo, grapeseed extract, green tea catechins, horsetail extract, L-cysteine, acetyl-L-carnitine, rosemary, saw palmetto, spirulina, probiotics, and various proprietary blends. Evidence quality across this category varies widely. Some, like marine collagen peptides and specific amino acid blends, appear in the multi-nutrient formulations discussed above, where controlled trials show measurable benefit as part of a combination product (Milani & Colombo, 2023). Others have plausible mechanistic rationale, such as antioxidant or anti-inflammatory activity, but lack controlled human trials needed to support specific clinical claims (Ahmed et al., 2025). The evidence-informed position is to treat this category as an area of active but immature research: reasonable to discuss with clients as adjunctive support, but not a substitute for confirming and correcting the core deficiencies outlined above, and not a category where strong efficacy claims are currently defensible.

Conclusion

The nutritional foundations mapped here are the raw materials the follicle needs, but raw materials require the right biological environment to become hair. Across protein, iron, the B-vitamins, zinc, selenium, the fat-soluble vitamins, and omega-3 fatty acids, the evidence converges on the same clinical pattern: deficiency is more prevalent among women with hair loss than conventional screening assumes (Almohanna et al., 2019; Gomes et al., 2025); conventional reference ranges frequently miss the functional thresholds the follicle actually requires (Al-Fawaeir & Al-Odat, 2025; Durusu Turkoglu et al., 2024); and gut absorptive capacity determines whether a well-designed nutrition or supplement plan can actually reach the tissue it is meant to support (Liu & Liu, 2023).

For nutrition professionals evaluating hair loss in practice, three principles should guide the work. First, test rather than guess: functional laboratory testing, not symptom-based assumptions, should drive every supplementation decision. Second, evaluate the whole pattern rather than a single nutrient in isolation, since the strongest evidence to date supports multi-nutrient approaches over any single compound (Drake et al., 2023; Milani & Colombo, 2023). Third, treat gut health as a prerequisite rather than an afterthought, since malabsorption can render an otherwise well-designed protocol ineffective. Hair loss in women is rarely explained by a single variable, but nutritional status is consistently among the most correctable factors and deserves a central place in how nutrition professionals evaluate and support clients navigating this condition.

References

Ahmed, A., Sharma, D., & Ungar, B. (2025). Impact of nutraceutical supplements on hair growth in non-scarring alopecia: A systematic review and meta-analysis. *Archives of Dermatological Research, 317*(1), 801. https://doi.org/10.1007/s00403-025-04306-8

Al-Fawaeir, S., & Al-Odat, I. (2025). Quantitative analysis of selected circulating hematological biomarkers, essential minerals, vitamins, and thyroid hormones in females affected by hair loss. *Diseases, 13*(11), 352. https://doi.org/10.3390/diseases13110352

Almohanna, H. M., Ahmed, A. A., Tsatalis, J. P., & Tosti, A. (2019). The role of vitamins and minerals in hair loss: A review. *Dermatology and Therapy, 9*(1), 51–70. https://doi.org/10.1007/s13555-018-0278-6

Beoy, L. A., Woei, W. J., & Hay, Y. K. (2010). Effects of tocotrienol supplementation on hair growth in human volunteers. *Tropical Life Sciences Research, 21*(2), 91–99.

Chang, H. C., & Chang, Y. S. (2022). Association between serum zinc levels and androgenetic alopecia: A systematic review and meta-analysis. *Journal of Cosmetic Dermatology, 21*(4), 1774–1777. https://doi.org/10.1111/jocd.14564

Drake, L., Reyes-Hadsall, S., Martinez, J., Heinrich, C., Huang, K., & Mostaghimi, A. (2023). Evaluation of the safety and effectiveness of nutritional supplements for treating hair loss: A systematic review. *JAMA Dermatology, 159*(1), 79–86. https://doi.org/10.1001/jamadermatol.2022.4867

Durusu Turkoglu, I. N., Turkoglu, A. K., Soylu, S., Gencer, G., & Duman, R. (2024). A comprehensive investigation of biochemical status in patients with telogen effluvium: Analysis of Hb, ferritin, vitamin B12, vitamin D, thyroid function tests, zinc, copper, biotin, and selenium levels. *Journal of Cosmetic Dermatology, 23*(12), 4277–4284. https://doi.org/10.1111/jocd.16512

Goette, D. K., & Odom, R. B. (1976). Alopecia in crash dieters. *JAMA, 235*(24), 2622–2623.

Gomes, N., Silva, N., & Teixeira, B. (2025). Assessing the relationship between dietary factors and hair health: A systematic review. *Nutrition and Health, 32*(4). https://doi.org/10.1177/02601060251367206

Liran, O., Vered, S., Cohen, B., Shapiro Ben David, S., Nakhleh, A., Rahamim-Cohen, D., Azuri, J., & Adler, L. (2025). Hair loss and zinc deficiency: A cross-sectional study. *Healthcare, 13*(22), 2965. https://doi.org/10.3390/healthcare13222965

Liu, Z., & Liu, X. (2023). Gut microbiome, metabolome and alopecia areata. *Frontiers in Microbiology, 14*, 1281660. https://doi.org/10.3389/fmicb.2023.1281660

Milani, M., & Colombo, F. (2023). Efficacy and tolerability of an oral supplement containing amino acids, iron, selenium, and marine hydrolyzed collagen in subjects with hair loss (androgenetic alopecia, AGA or FAGA or telogen effluvium): A prospective, randomized, 3-month, controlled, assessor-blinded study. *Skin Research and Technology, 29*(6), e13381. https://doi.org/10.1111/srt.13381

Moltó-Balado, P., Simeó-Monzo, A., & del Barrio-Gonzalez, A. (2026). Effectiveness of biotin supplementation for hair growth in patients with alopecia: A systematic review. *Dermato, 6*(2), 17. https://doi.org/10.3390/dermato6020017

Rangarajan, S. (2025). Assessment of serum ferritin levels in female patients with telogen effluvium. *Cureus, 17*(12), e100249. https://doi.org/10.7759/cureus.100249

Suo, L., VanBuren, C., Hovland, E. D., Kedishvili, N. Y., Sundberg, J. P., & Everts, H. B. (2021). Dietary vitamin A impacts refractory telogen. *Frontiers in Cell and Developmental Biology, 9*, 571474. https://doi.org/10.3389/fcell.2021.571474

Yongpisarn, T., Tejapira, K., Thadanipon, K., & Suchonwanit, P. (2024). Vitamin D deficiency in non-scarring and scarring alopecias: A systematic review and meta-analysis. *Frontiers in Nutrition, 11*, 1479337. https://doi.org/10.3389/fnut.2024.1479337

About the Author

Julie Olson, BCHN®, CN, CFMP, CGP, BSc, is a Board-Certified Holistic Nutritionist and the founder of Fortitude Functional Nutrition and the HealthyHairFix® program. She works with women experiencing hair thinning and loss, also known as Alopecia, using functional lab testing and root-cause nutrition strategies to identify and correct deficiencies that conventional workups often miss.